

Stop Taking These Medications Prior To Testing

It is **important** to **stop** all medications listed or your **appointment** will need to be **canceled**.



ANTIHISTAMINES

Stop taking these 7-10 days prior to testing

Allegra (Fexofenadine) Claritin (Loratadine) Zyrtec (Cetirizine) Xyzal

Stop taking these 72 Hours prior to testing (3 days)

Benadryl
Dymista
Tylenol Allergy Sinus
Tylenol Cold & Flu
Tylenol Flu PM
Azelastine
Doxylamine
Olopatadine

PLEASE CONTINUE
TO TAKE ALL INHALERS
AND NASAL SPRAYS
AS DIRECTED BY
YOUR PROVIDER



SLEEP AIDS

Stop taking these 72 Hours prior to testing (3 days)

Tylenol PM Unisom



BLOCKING MEDICATIONS

Do not take day of testing Singular (Montelukast)

Zyfo (Zileuon) Accolate (Zafirlukast)



ANTACID MEDICATIONS

Stop taking these 24 Hours prior to testing (1 day)

Ok to take Tums or Rolaids if needed Pepcid (Famotidine) Zantac (Ranitidine)

Tagamet (Cimetidine)



ORAL STEROIDS

Stop taking 2 weeks prior to testing

Prednisone

TOPICAL STEROIDS

at the site of testing

paraccess.com | 804-320-4243

What To Expect During Your Allergy Testing?



Testing process takes about 45 minutes depending on your — reactions. —

Testing is done on the back. Once allergens have been placed, they

have to sit for 15 minutes.

we test for an

expanded environmental panel—
animals, trees, grasses, mold
and weeds— and a food panel
(as ordered by the provider.)





If you have reactions to any of the allergens tested it will raise up, turn red and Itch.



- We do not use needles. We use - scratchers; it's just enough to get a antigen into the surface of the skin.

If an area itches

DO NOT
scratch the area.

WE CAN NOT TEST YOU IF YOU'VE HAD OR HAVE ANY OF THE FOLLOWING:



Pregnant



Tattoos (that cover the back area)



Rash or Sunburn (on back area)

